

AD\_\_\_\_\_

Award Number: W81XWH-11-1-0452

TITLE: Intranasal Oxytocin for the Treatment of Pain Associated with Interstitial Cystitis

PRINCIPAL INVESTIGATOR: Meredith T. Robbins, Ph.D.

CONTRACTING ORGANIZATION: University of Alabama at Birmingham  
Birmingham, AL 35294

REPORT DATE: November 2015

TYPE OF REPORT: Final

PREPARED FOR: U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

DISTRIBUTION STATEMENT: Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.					
1. REPORT DATE November 2015		2. REPORT TYPE Final		3. DATES COVERED 1 Sep 2011 - 31 Aug 2015	
4. TITLE AND SUBTITLE  Intranasal Oxytocin for the Treatment of Pain Associated with Interstitial Cystitis				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-11-1-0452	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Meredith T. Robbins, Ph.D.  E-Mail: mturnbach@yahoo.com				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  University of Alabama at Birmingham 701 20 <sup>th</sup> St. South Birmingham, AL 35294-0001				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT The purpose of this study was to evaluate the efficacy of intranasal oxytocin on bladder nociception in a cohort of patients with interstitial cystitis (IC). We hypothesized that patients with IC are more likely to experience treatment responses as defined by global response assessment (GRA) scores when they receive intranasal oxytocin as opposed to when they receive intranasal saline. Secondary outcome measures included a verbal pain report (VPR; 0-10 with 0 being no pain and 10 being the worst possible pain), a verbal anxiety report (VAR; 0-10 with 0 being no anxiety and 10 being the worst possible anxiety), number of voids since last contact with an investigator, and interim medications used for pain control or anxiety. Compared to saline administration intranasal oxytocin did not significantly decrease anxiety, pain or the number of voids reported by participants.					
15. SUBJECT TERMS Interstitial cystitis, oxytocin, intranasal, bladder, pain, anxiety					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU		19b. TELEPHONE NUMBER (include area code)

## Table of Contents

	<u>Page</u>
1. Introduction.....	4
2. Keywords.....	4
3. Accomplishments.....	4-8
4. Impact.....	8-9
5. Changes/Problems.....	9
6. Products.....	9-10
7. Participants & Other Collaborating Organizations.....	10
8. Special Reporting Requirements.....	10
9. Appendices.....	10
10. References.....	10-11

## 1. INTRODUCTION

The purpose of this study was to evaluate the efficacy of intranasal oxytocin on bladder nociception in a cohort of patients with interstitial cystitis (IC). We hypothesized that patients with IC would be more likely to experience treatment responses as defined by global response assessment (GRA) scores when they received intranasal oxytocin as opposed to when they received intranasal saline. Our primary outcome measure was the GRA score, which was collected at 6 and 24 hours post drug or placebo administration. This is a seven-point symmetric scale previously validated for use in IC studies in which patients are asked relative to baseline (i.e. over the last 6 hours for purposes of this study), are you markedly worse, moderately worse, slightly worse, no change, slightly improved, moderately improved, or markedly improved. Moderate or marked improvement was defined as a treatment response. Secondary outcome measures included a verbal pain report (VPR; 0-10 with 0 being no pain and 10 being the worst possible pain), a verbal anxiety report (VAR; 0-10 with 0 being no anxiety and 10 being the worst possible anxiety), number of voids since last contact with an investigator, and interim medications used for pain control or anxiety. The VPR and VAR were assessed for the current point in time and the interim time period since last investigator-initiated contact. Participants were asked to record their number of voids and concomitant medication use in a diary so that the results could be accurately reported to the investigators.

## 2. KEYWORDS

Interstitial cystitis  
Oxytocin  
Bladder  
Global response assessment  
Verbal pain report  
Verbal anxiety report  
Void

## 3. ACCOMPLISHMENTS

### ▪ What were the major goals of the project?

The major goal of this project was to evaluate the efficacy of intranasal oxytocin in relieving bladder pain in a cohort of patients with interstitial cystitis.

### ▪ What was accomplished under these goals?


Our primary outcome measure was the global response assessment (GRA) score, which was collected at 6 and 24 hours post drug or placebo administration. This is a seven-point symmetric scale previously validated for use in IC studies in which patients are asked relative to baseline (over the last 6 hours for purposes of this study), are you markedly worse, moderately worse, slightly worse, no change, slightly improved, moderately improved, or markedly improved. Moderate or marked improvement will be defined as a treatment response.<sup>1-2</sup>

Secondary outcome measures included a verbal pain report (VPR; 0-10 with 0 being no pain and 10 being the worst possible pain), a verbal anxiety report (VAR; 0-10 with 0 being no anxiety and 10 being the worst possible anxiety), number of voids since last contact with an


investigator, and interim medications used for pain control or anxiety. The VPR and VAR were assessed for the current point in time and the interim time period since last investigator-initiated contact.

For purposes of comparison with our previous studies and those of others related to subjects with pain, we administered a series of questionnaires that characterized the subjects' demographic information, health history, pain-related symptoms, health status perceptions and hypervigilance.


Figure 1 below shows mean GRA scores obtained 6 and 24 hours post oxytocin or saline administration. There was no significant difference in GRA scores obtained after patients received oxytocin compared to when they received saline. Furthermore, neither oxytocin nor saline produced moderate or marked improvement, which would have defined a treatment response.



Secondary response measures that were collected and analyzed for possible treatment effects were the VPR, VAR and number of voids. In Figure 2 below, it is clear that both the VPR (left panel) and the VAR (right panel) both decreased after patients intranasally administered oxytocin and saline. However, there was no significant difference in improvement in these measures post-oxytocin compared to post-saline. Similarly, voiding behavior was not different after intranasal oxytocin compared to intranasal saline. Figure 3 shows total number of voids in the first 6 hours (left panel) and over the entire 24 hour (right panel) evaluation period.



**Figure 2: Secondary Outcome Measures.** VPR (left panel) and VAR (right panel) were assessed 1, 2, 4, 6 and 24 hours after oxytocin and saline administration. Although pain and anxiety reported by patients did decrease, this effect was apparent regardless of the type of intranasal spray administered. No significant differences were found (N=25).



**Figure 3: Voiding Behavior.** The total number of voids in the first 6 h and over the entire 24 h experimental period was calculated. Patients' voiding was similar at both time points regardless of the intranasal substance received, and no significant differences were observed over the first 6 h ( $t=0.327$ ;  $p=0.745$ ) nor over 24 h ( $t=0.980$ ;  $p=0.333$ ). N=23-24.

General demographic and health history information were gathered using a traditional medical history and custom form. The O'Leary-Sant IC Symptom and Problem Index<sup>3</sup> was utilized to assess urinary symptoms. The Center for Epidemiologic Studies - Depression Scale (CES-D) was administered to assess symptoms of depression.<sup>4-5</sup> The McGill Pain Questionnaire (MPQ), a self-report instrument that measures the sensory, affective, and intensity dimensions of pain, was also administered.<sup>6-8</sup> These data are shown in Tables 1 and 2.

**Table 1. Characteristics of the Participants in the Study Sample**

Characteristic	
Age (mean $\pm$ SD)	45.80 $\pm$ 12.63
Sex (n, %)	
Women	25 (100)
Men	0 (0)
Ethnic background (n, %)	
Caucasion	23 (92)
African-American	1 (4)
Native-American	1 (4)
Education in years (mean $\pm$ SD)	13.84 $\pm$ 1.99
Comorbidities (n, %)	
TMJ/Facial Pain	9 (36)
Fibromyalgia	8 (32)
Chronic Fatigue Syndrome	9 (36)
Back Pain	13 (52)
Joint Pain	9 (36)
Chronic Headache	8 (32)
Depression	12 (48)
IBS/ GI Pain	10 (40)
Premenstrual/Menstrual Pain	14 (56)
Bladder Pain	25 (100)

**Table 2. Questionnaire Data from the Participants in the Study Sample**

Questionnaire	Score (mean $\pm$ SD)
Short-Form McGill Pain Questionnaire	
Sensory Pain Rating Index (range 0-33)	15.12 $\pm$ 5.90
Affective Pain Rating Index (range 0-12)	5.40 $\pm$ 3.12
Total Pain Rating Index	20.52 $\pm$ 8.51
Evaluative Overall Intensity of Pain Experience (range 0-5)	3.04 $\pm$ 0.88
IC Symptom and Problem Questionnaire	
IC Symptom Index (range 0-20)	13.36 $\pm$ 4.95
IC Problem Index (range 0-16)	10.92 $\pm$ 4.06
Center for Epidemiologic Studies Depression Scale (range 0-60)	23.96 $\pm$ 6.57

- **What opportunities for training and professional development has the project provided?**

This project allowed me the opportunity to design and execute a clinical trial. This was an invaluable experience. Furthermore, I was able to interact with a population of patients with chronic pain for the first time. Up to this point in my career I had only used animal models to study pain in general and interstitial cystitis specifically. To be able to speak with patients who live with this condition was a truly eye-opening experience. I gained a much better understanding of the disease itself and of chronic pain while working on this project.

- **How were the results disseminated to communities of interest?**

Nothing to Report

- **What do you plan to do during the next reporting period to accomplish the goals?**

Nothing to Report

#### **4. IMPACT**

- **What was the impact on the development of the principal discipline(s) of the project?**

Unfortunately there was no apparent beneficial effect of the intranasal oxytocin in the present study. It may be that the current delivery system was ineffective in allowing the entire dose of oxytocin to reach its target and/or be absorbed completely. The volume necessary to deliver the appropriate dose of oxytocin is quite high, so it is possible that switching to a higher concentration/lower volume formulation of oxytocin would be more efficient and would allow for definitive determination of its effectiveness at relieving pain and/or anxiety.

- **What was the impact on other disciplines?**

Nothing to Report

- **What was the impact on technology transfer?**

Nothing to Report

- **What was the impact on society beyond science and technology?**

Nothing to Report

## **5. CHANGES/PROBLEMS**

- **Changes in approach and reasons for change**

Nothing to Report

- **Actual or anticipated problems or delays and actions or plans to resolve them**

Recruitment was more difficult than initially anticipated. For this reason, a no-cost extension request for an additional 12 months of time was submitted, which allowed for completion of the study as originally proposed. No additional funds were requested for this time.

- **Changes that had a significant impact on expenditures**

Nothing to Report

- **Significant changes in use or care of human subjects, vertebrate animals, biohazards, and/or select agents**

Nothing to Report

## **6. PRODUCTS**

- **Publications, conference papers, and presentations**

- **Journal publications.** Nothing to Report

- **Books or other non-periodical, one-time publications.** Nothing to Report

- **Other publications, conference papers, and presentations.** Nothing to Report

- **Website(s) or other Internet site(s)**

- <https://clinicaltrials.gov/>

- **Technologies or techniques**

Nothing to Report

- **Inventions, patent applications, and/or licenses**

Nothing to Report

- **Other Products**

Nothing to Report

## **7. PARTICIPANTS & OTHER COLLABORATING ORGANIZATIONS**

- **What individuals have worked on the project?**

Meredith T. Robbins-no change

Timothy J. Ness-no change

Michael Froelich-no change

L. Keith Lloyd-no change

- **Has there been a change in the active other support of the PD/PI(s) or senior/key personnel since the last reporting period?**

- The PI, Meredith Robbins, has closed two previously active NIH grants. One study was completed in June 2015, and the second study was completed in September 2015.

- **What other organizations were involved as partners?**

Nothing to Report

## **8. SPECIAL REPORTING REQUIREMENTS**

N/A

## **9. APPENDICES**

N/A

## **10. REFERENCES**

1. Probert KJ, Mayer RD, Wang Y, Sant GR, Hanno PM, Peters KM, Kusek JW; Interstitial Cystitis Clinical Trials Group: Responsiveness of symptom scales for interstitial cystitis. Urology 67(1): 55-59, 2006.

2. Nickel JC, Moldwin R, Lee S, Davis EL, Henry RA, Wyllie MG: Intravesical alkalinized lidocaine (PSD597) offers sustained relief from symptoms of interstitial cystitis and painful bladder syndrome. *BJU Int* 103(7): 910-18, 2009.
3. O'Leary MP, Sant GR, Fowler FJ, Whitmore KE, Spolarich-Kroll J: The interstitial cystitis symptom index and problem index. *Urology*, suppl 49: 58, 1997.
4. Radloff L: The CES-D Scale: A self-report depression scale for research in the general population, *Appl Psychol Measurement* 1; 385-401, 1977.
5. Bradley LA, Richter JE, Pulliam TJ, McDonald-Haile J., Scarinci IC, Schan CA, Dalton CB and Salley AN: The relationship between stress and symptoms of gastroesophageal reflux: the influence of psychological factors. *Am J Gastroenterol.* 88: 11-19, 1993.
6. Melzack R: The McGill Pain Questionnaire: Major properties and scoring methods. *Pain* 1: 277-299, 1975.
7. Melzack R (ed.): *Pain measurement and assessment*. New York; Raven Press, 1983.
8. Melzack R: The Short-Form McGill Pain Questionnaire. *Pain*, 30: 191-197, 1987.